

Please click price below and you will be redirected to our pharmacy.

## Ciloxan



Generic **Ciloxan** is used for treating eye infections caused by certain bacteria.

Active ingredient: [Ciprofloxacin](#)

**Brand(s):** Ciplox Eye Drops / Ziflox

**Manufacturer:** Cipla Limited / Zen Pharma (P) Ltd.

**Disease(s):** Infections

### Ciloxan 0.3% 5ml

|            | Per solution | Per Pack                              | Order                       |
|------------|--------------|---------------------------------------|-----------------------------|
| 1 solution | \$ 12.00     | <b>\$ 12.00</b>                       | <a href="#">Add to cart</a> |
| 2 solution | \$ 11.00     | <b>\$ 22.00</b><br><del>\$24.00</del> | <a href="#">Add to cart</a> |
| 3 solution | \$ 10.00     | <b>\$ 30.00</b><br><del>\$36.00</del> | <a href="#">Add to cart</a> |
| 4 solution | \$ 9.00      | <b>\$ 36.00</b><br><del>\$48.00</del> | <a href="#">Add to cart</a> |
| 5 solution | \$ 8.00      | <b>\$ 40.00</b><br><del>\$60.00</del> | <a href="#">Add to cart</a> |
| 6 solution | \$ 7.00      | <b>\$ 42.00</b><br><del>\$72.00</del> | <a href="#">Add to cart</a> |

## Ciprofloxacin 0.3% solution

### What is this medicine?

CIPROFLOXACIN is a quinolone antibiotic. It is used to treat bacterial eye infections.

### What should my health care professional know before I take this medicine?

They need to know if you have any of these conditions:

- contact lens wearer
- an unusual or allergic reaction to ciprofloxacin, other antibiotics or medicines, foods, dyes, or preservatives
- pregnant or trying to get pregnant
- breast-feeding

### How should I take this medicine?

This medicine is only for use in the eye. Wash hands before and after use. Try not to touch the tip of the dropper to anything, even your eye or fingertips. Tilt your head back slightly and pull your lower eyelid down with your index finger to form a pouch. Squeeze the prescribed number of drops into the pouch. Close the eye gently to spread the drops. Your vision may blur for a few minutes. Use your doses at regular intervals. Do not use your medicine more often than directed. Finish the full course that is prescribed even if you think your condition is better. Do not skip doses or stop your medicine early.

Talk to your pediatrician regarding the use of this medicine in children. Special care may be needed.

Overdosage: If you think you have taken too much of this medicine contact a poison control center or emergency room at once.

