

Proair



ProAir is used to treat wheezing and shortness of breath caused by breathing problems (such as asthma, chronic obstructive pulmonary disease). ProAir belongs to a class of drugs known as bronchodilators. ProAir works in the airways by opening breathing passages and relaxing muscles.

Active ingredient: [Albuterol \(salbutamol\)](#)

Brand(s):Asthalin

Manufacturer: Cipla

Disease(s):Asthma / Bronchospasm

Proair 100mcg

	Per item	Per Pack	Order
1 inhaler	\$ 22.95	\$ 22.95	Add to cart
2 inhaler	\$ 19.98	\$ 39.95 \$45.90	Add to cart
3 inhaler	\$ 18.32	\$ 54.95 \$68.85	Add to cart
4 inhaler	\$ 17.49	\$ 69.95 \$91.80	Add to cart
6 inhaler	\$ 16.66	\$ 99.95 \$137.70	Add to cart

Albuterol Inhaler

What is this medicine?

Albuterol(also known as salbutamol) is a bronchodilator that relaxes muscles in the airways and increases air flow to the lungs. Albuterol inhalation is used to treat or prevent bronchospasm in people with reversible obstructive airway disease. Albuterol is also used to prevent exercise-induced bronchospasm. Albuterol inhalation may also be used for purposes not listed in this medication guide.

How should I use this medicine?

Use albuterol inhalation exactly as prescribed by your doctor. Do not use in larger or smaller amounts or for longer than recommended. Follow the directions on your prescription label.

Use only the prescribed dose of this medicine and follow all patient instructions for safe use. Talk with your doctor about your individual risks and benefits of using albuterol.

When using the albuterol inhaler device for the first time, prime it by spraying 4 test sprays into the air, away from your face. Shake well before priming. Also prime the inhaler if you have not used it for 2 weeks or longer, or if you have dropped the inhaler.

The instructions below are for standard use of the inhaler and nebulizer devices. Your doctor may want you to use your device differently. Use only the inhaler device provided with your medicine or you may not get the correct dose.

To use the albuterol inhaler:

- Shake the canister well just before each spray.
- Uncap the mouthpiece of the albuterol inhaler. Breathe out fully. Put the mouthpiece into your mouth and close your lips. Breathe in slowly while pushing down on the canister. Hold your breath for 10 seconds, then breathe out slowly.
- If you use more than one inhalation at a time, wait at least 1 minute before using the second inhalation and shake the inhaler again.

