

Please click price below and you will be redirected to our pharmacy.

Procoralan



Procoralan works by affecting your heart's electrical activity in order to slow the heart rate. Ivabradine is used in certain people with chronic heart failure, to help lower the risk of needing to be hospitalized when symptoms get worse. Ivabradine is not for use if you already have heart failure symptoms that recently got worse before starting this medicine. Ivabradine may also be used for purposes not listed in this medication guide.

Active ingredient: [Ivabradine](#)

Brand(s):Ivabeat

Manufacturer: Cipla Limited

Disease(s):Chronic Heart Failure

Procoralan 5mg

	Per pill	Per Pack	Order
10 tablet	\$ 2.50	\$ 24.95	Add to cart
20 tablet	\$ 2.00	\$ 39.95 \$49.90	Add to cart
30 tablet	\$ 1.67	\$ 49.95 \$74.85	Add to cart
60 tablet	\$ 1.50	\$ 89.95 \$149.70	Add to cart
90 tablet	\$ 1.44	\$ 129.95 \$224.55	Add to cart
120 tablet	\$ 1.37	\$ 164.95 \$299.40	Add to cart
Free AirMail shipping			
180 tablet	\$ 1.33	\$ 239.95 \$449.10	Add to cart
Free AirMail shipping			

Ivabradine tablet

What is this medicine?

IVABRDINE works by affecting your heart's electrical activity in order to slow the heart rate. Ivabradine is used in certain people with chronic heart failure, to help lower the risk of needing to be hospitalized when symptoms get worse. Ivabradine is not for use if you already have heart failure symptoms that recently got worse before starting this medicine. Ivabradine may also be used for purposes not listed in this medication guide.

What should I tell my health care providers before I take this medicine?

You should not use ivabradine if you are allergic to it, or if you have:

- a serious heart condition such as "sick sinus syndrome" or "AV block" (unless you have a pacemaker);
- heart failure symptoms that recently got worse;
- severe liver disease;
- if your resting heart rate is less than 60 beats per minute;
- if your blood pressure is lower than 90 over 50; or
- if you depend on a pacemaker to regulate your heart rate.

Some medicines can cause unwanted or dangerous effects when used with ivabradine. Tell your doctor if you are taking any other medicines, including over-the-counter medicines, vitamins, or herbal products.

