

Please click price below and you will be redirected to our pharmacy.

Skelaxin



Generic Skelaxin is a muscle relaxant. It works by blocking nerve impulses (or pain sensations) in the brain. **Generic Skelaxin** is used together with rest and physical therapy to treat discomfort associated with acute skeletal muscle conditions such as pain or injury.

Active ingredient: [Metaxalone Mr](#)

Brand(s): Flexura

Manufacturer: Sun Pharmaceutical Industries Ltd.

Disease(s): Acute Skeletal Muscle Condition / Pain / Injury

Skelaxin 400mg

| | Per pill | Per Pack | Order |
|-----------------------|----------|---|-----------------------------|
| 10 tablet | \$ 1.99 | \$ 19.95 | Add to cart |
| 20 tablet | \$ 2.25 | \$ 44.95 \$44.95 | Add to cart |
| 30 tablet | \$ 2.67 | \$ 79.95 \$79.95 | Add to cart |
| 60 tablet | \$ 1.83 | \$ 109.95 \$119.70 | Add to cart |
| 90 tablet | \$ 1.61 | \$ 144.95 \$179.55 | Add to cart |
| 120 tablet | \$ 1.58 | \$ 189.95 \$239.40 | Add to cart |
| Free AirMail shipping | | | |
| 180 tablet | \$ 1.00 | \$ 179.95 \$359.10 | Add to cart |

Free AirMail shipping

Metalaxone tablet

What is this medicine?

METAXALONE is a muscle relaxant. It works by blocking nerve impulses (or pain sensations) in the brain.

Metalaxone is used together with rest and physical therapy to treat discomfort associated with acute skeletal muscle conditions such as pain or injury.

What should I tell my health care provider before I take this medicine?

You should not use Metalaxone if you are allergic to metaxalone, or if you have:

- anemia (a lack of red blood cells);
- severe kidney disease; or
- severe liver disease.

Before using Metalaxone, tell your doctor if you are allergic to any drugs, or if you have liver or kidney disease. You may need a dose adjustment or special tests to safely take this medicine.

Metalaxone may be harmful to an unborn baby. Tell your doctor if you are pregnant or plan to become pregnant during treatment. It is not known whether metaxalone passes into breast milk or if it could harm a nursing baby. Do not use this medication without telling your doctor if you are breast-feeding a baby.

