

Please click price below and you will be redirected to our pharmacy.

## Diamox



Generic **Diamox** is used for treating certain types of glaucoma in combination with other medicines. It is also used to treat or prevent symptoms of mountain sickness.

Active ingredient: [Acetazolamide](#)

**Brand(s):**Avva

**Manufacturer:** Intas Pharmaceuticals Ltd.

**Disease(s):**Glaucoma / Mountain Sickness

### Diamox 250mg

|            | Per pill | Per Pack                                | Order                       |
|------------|----------|---|-----------------------------|
| 30 tablet  | \$ 1.00  | <b>\$ 29.95</b>                         | <a href="#">Add to cart</a> |
| 60 tablet  | \$ 0.93  | <b>\$ 55.95</b><br><del>\$59.90</del>   | <a href="#">Add to cart</a> |
| 90 tablet  | \$ 0.89  | <b>\$ 79.95</b><br><del>\$89.85</del>   | <a href="#">Add to cart</a> |
| 120 tablet | \$ 0.83  | <b>\$ 99.95</b><br><del>\$119.80</del>  | <a href="#">Add to cart</a> |
| 180 tablet | \$ 0.72  | <b>\$ 129.95</b><br><del>\$179.70</del> | <a href="#">Add to cart</a> |
| 360 tablet | \$ 0.69  | <b>\$ 249.95</b><br><del>\$359.40</del> | <a href="#">Add to cart</a> |

Free AirMail shipping

## Acetazolamide tablet

### What is this medicine?

ACETAZOLAMIDE is used to treat glaucoma and some seizure disorders. It may be used to treat edema or swelling from heart failure or from other medicines. This medicine is also used to treat and to prevent altitude or mountain sickness.

### What should I tell my health care provider before I take this medicine?

They need to know if you have any of these conditions:

- diabetes
- kidney disease
- liver disease
- lung disease
- an unusual or allergic reaction to acetazolamide, sulfa drugs, other medicines, foods, dyes, or preservatives
- pregnant or trying to get pregnant
- breast-feeding

### How should I use this medicine?

Take this medicine by mouth with a glass of water. Take this medicine with food if it upsets your stomach. Take your doses at regular intervals. Do not take your medicine more often than directed. Do not stop taking except on your doctor's advice.

Talk to your pediatrician regarding the use of this medicine in children. Special care may be needed.

Patients over 65 years old may have a stronger reaction and need a smaller dose.

