

## Ticlid



Generic **Ticlid** is used for reducing the risk of stroke. It may also be used in combination with aspirin to prevent a blood clot from forming after a stent is placed in the coronary artery.

Active ingredient: **Ticlopidine**

**Brand(s):**Ticlop

**Manufacturer:** Zydus Cadila Healthcare

**Disease(s):**Stroke / Blood Clots

### Ticlid 250mg

	Per pill	Per Pack	Order
30 tablet	\$ 2.33	<b>\$ 69.95</b>	<a href="#">Add to cart</a>
60 tablet	\$ 2.17	<b>\$ 129.95</b> <del>\$139.90</del>	<a href="#">Add to cart</a>
90 tablet	\$ 2.11	<b>\$ 189.95</b> <del>\$209.85</del>	<a href="#">Add to cart</a>
Free AirMail shipping			
120 tablet	\$ 2.08	<b>\$ 249.95</b> <del>\$279.80</del>	<a href="#">Add to cart</a>
Free AirMail shipping			
180 tablet	\$ 2.09	<b>\$ 376.80</b> <del>\$419.70</del>	<a href="#">Add to cart</a>

Free AirMail shipping

## Ticlopidine tablet

### What is this medicine?

TICLOPIDINE helps to prevent blood clots. This medicine is used to prevent strokes in patients who have already had a stroke, or those who are at high risk of having a stroke. This medicine may also be used in combination with aspirin to prevent blood clots in patients who have had a coronary stent placed.

### What should my health care professional know before I take this medicine?

They need to know if you have any of the following conditions:

- anemia
- high cholesterol
- history of bleeding disorder
- kidney disease
- liver disease
- stomach ulcer
- an unusual or allergic reaction to ticlopidine, other medicines, foods, dyes, or preservatives
- pregnant or trying to get pregnant
- breast-feeding

### How should I take this medicine?

Take this medicine by mouth with a glass of water. Take with food or just after eating to reduce stomach upset. Take your doses at regular intervals. Do not take your medicine more often than directed.

