

## Tobrex



**Tobrex** (for the eyes) is used to treat bacterial infections of the eyes. **Tobrex** will not treat a viral or fungal infection of the eye. **Tobrex** is for use in treating only bacterial infections.

Active ingredient: [Tobramycin](#)

**Brand(s):**Tobastar / Toba

**Manufacturer:** Lifestar / Sun Pharma

**Disease(s):**Bacterial Infections Of The Eyes

### Tobrex 0.3% 5ml

	Per pill	Per Pack	Order
1 dropper	\$ 29.95	<b>\$ 29.95</b>	<a href="#">Add to cart</a>
2 dropper	\$ 26.98	<b>\$ 53.95</b> <del>\$59.90</del>	<a href="#">Add to cart</a>
3 dropper	\$ 21.65	<b>\$ 64.95</b> <del>\$89.85</del>	<a href="#">Add to cart</a>
4 dropper	\$ 18.24	<b>\$ 72.95</b> <del>\$119.80</del>	<a href="#">Add to cart</a>
5 dropper	\$ 15.99	<b>\$ 79.95</b> <del>\$149.75</del>	<a href="#">Add to cart</a>

## Tobrex Eye Drops

### What is tobramycin inhalation?

Tobramycin is an antibiotic that is inhaled into the lungs using a nebulizer. Tobramycin inhalation is used to treat lung infections in patients with cystic fibrosis.

tobramycin is for use in adults and children who are at least 6 years old. Tobramycin inhalation may also be used for purposes not listed in this medication guide.

### Important Information

Follow all directions on your medicine label and package. Tell each of your healthcare providers about all your medical conditions, allergies, and all medicines you use.

### Before taking this medicine

You should not use tobramycin if you are allergic to tobramycin or similar antibiotics (amikacin, gentamicin, kanamycin, neomycin, paromomycin, streptomycin).

Tell your doctor if you have ever had:

- hearing problems;
- breathing problems (cough, wheezing, chest tightness);
- dizzy spells;
- a neuromuscular disease such as myasthenia gravis or Parkinson's disease; or
- kidney disease.

It is not known whether tobramycin inhalation will harm an unborn baby. However, a medicine similar to tobramycin caused irreversible hearing damage in babies whose mothers took that medicine during pregnancy. Tell your doctor if you are pregnant or plan to become pregnant. Having cystic fibrosis during pregnancy may increase the risk of premature birth. The benefit of treating cystic fibrosis may outweigh any risks to the baby. If you breastfeed while using tobramycin, watch the baby for signs of diarrhea, blood in the stools, or a fungal diaper rash with white patches on the skin.

