

Please click price below and you will be redirected to our pharmacy.

## Tobradex



**Tobradex** ophthalmic is used to treat bacterial infections of the eyes. Tobramycin is an antibiotic. It is used to treat bacterial infections. Dexamethasone is a steroid. Dexamethasone ophthalmic is used to treat the swelling associated with bacterial infections of the eye.

Active ingredient: [Tobramycin/dexamethasone](#)

**Brand(s):**Toba DM

**Manufacturer:** Sun Pharma

**Disease(s):**Swelling / Bacterial Infections

### Tobradex 0.1%/0.3% 10ml

	Per pill	Per Pack	Order
1 dropper	\$ 9.95	<b>\$ 9.95</b>	<a href="#">Add to cart</a>
2 dropper	\$ 7.47	<b>\$ 14.95</b> <del>\$19.90</del>	<a href="#">Add to cart</a>
3 dropper	\$ 6.32	<b>\$ 18.95</b> <del>\$29.85</del>	<a href="#">Add to cart</a>
4 dropper	\$ 5.49	<b>\$ 21.95</b> <del>\$39.80</del>	<a href="#">Add to cart</a>
5 dropper	\$ 4.99	<b>\$ 24.95</b> <del>\$49.75</del>	<a href="#">Add to cart</a>

## Tobradex ophthalmic

### What is Tobramycin and Dexamethasone ophthalmic?

Tobradex ophthalmic contains a combination of tobramycin and dexamethasone. Tobramycin is an antibiotic. It is used to treat bacterial infections. Dexamethasone is a steroid. Dexamethasone ophthalmic is used to treat the swelling associated with bacterial infections of the eye.

Tobramycin and Dexamethasone ophthalmic is used to treat bacterial infections or inflammation of the eyes. It is available as an ointment and a liquid suspension.

### Important information

Contact your doctor if the symptoms begin to get worse or if you do not see any improvement in the condition after a few days.

Do not touch the dropper or tube opening to any surface, including your eyes or hands. The dropper or tube opening is sterile. If it becomes contaminated, it could cause an infection in the eye.

Apply light pressure to the inside corner of your eye (near your nose) after each drop to prevent the fluid from draining down your tear duct.

### What should I discuss with my doctor before using?

Do not use Tobramycin and Dexamethasone ophthalmic if you have a viral or fungal infection in the eye, such as herpes, vaccinia, or varicella. It is used to treat infections caused by bacteria only. Do not use Tobramycin and Dexamethasone ophthalmic for mycobacterial infections of the eye.

Tell your doctor if you are allergic to any aminoglycoside antibiotics (including gentamicin, streptomycin, tobramycin, others). This may cause an allergic reaction to Tobramycin and Dexamethasone ophthalmic.

It is not known whether Tobramycin and Dexamethasone ophthalmic will be harmful to an unborn baby. Do not use Tobramycin and Dexamethasone ophthalmic without first talking to your doctor if you are pregnant. It is not known whether tobramycin or dexamethasone passes into breast milk. Do not use this medication without first talking to your doctor if you are breast-feeding a baby.

### How should I use Tobramycin and Dexamethasone ophthalmic?

